

**North Central Research Extension Center—Minot
Dry Pea Variety Trial—Benson County**

Variety	Days to Maturity	Canopy		Disease 0-9	Lodging 0-9	Seed Weight g/1000	Test Weight lb/bu	Protein %	Seed Yield		
		Ht at Harvest in							2004 ----bu/A-----	2 Year	3 Year
CDC Mozart	101.8	12.7	0.0	6.8	251.2	64.2	19.4	98	--	--	
Carneval	97.8	24.8	2.8	2.8	189.8	65	19.2	87.9	--	--	
DS Admiral	99.8	23.2	0	4.8	226	65.2	18.8	92.6	--	--	
Majoret	100.8	19	2.5	5.8	207.7	64.8	19.9	86.1	--	--	
Cruiser	98.8	22.5	3.8	3.8	176.5	64.3	19.5	77	--	--	
Nitouche	98.5	22.7	5.5	3.5	245.7	64.1	19.7	88.6	--	--	
Stirling	102.8	8.2	0	7.8	200.4	63.5	21.9	73.9	--	--	
LSD 5%	1.79	4.6	0.9	1.8	15.5	0.6	1.2	12.9	--	--	
C.V.%	1.2	16.3	29.6	24.1	3	0.6	4	10	--	--	
Mean	100	19	2.1	5	213.9	64.4	19.8	86.3	--	--	

Disease score based on scale 0-9 (0=no powdery mildew)

Lodging score based on scale 0-9 (0=upright, 9=flat)

No data for 2002 or 2003, first year in rotation of counties.